

## **REAL-PE Study Demonstrates Statistically Significant Lower Major Bleeding Rates with the EKOS™ Endovascular System Compared to Mechanical Thrombectomy Device for Treatment of Pulmonary Embolism**

*Analysis is first to compare health system-based electronic health record data for treatment of pulmonary embolism with interventional devices*

MARLBOROUGH, Mass., Oct. 24, 2023 /PRNewswire/ -- Data from the REAL-PE study was presented today at Transcatheter Cardiovascular Therapeutics (TCT), the annual scientific symposium of the Cardiovascular Research Foundation, demonstrating that patients treated for pulmonary embolism (PE) with the Boston Scientific (NYSE: BSX) EKOS™ Endovascular System (EKOS) had lower rates of adverse events, including statistically significant lower rates of major bleeding, within seven days following their procedure compared to the Inari FlowTrieve® System. The analysis is the largest comparative study to use near real-time health system-based electronic health record (EHR) data to understand clinical practices and outcomes related to PE.

Each year, approximately 350,000 patients in the United States are affected by PE, a blood clot causing a blockage in one or more pulmonary arteries in the lungs and a leading cause of in-hospital death in the U.S. <sup>i,ii</sup> While treatment for PE has historically included anticoagulant medication, the use of new interventional therapies, such as the EKOS system, is increasing. <sup>iii</sup> The EKOS system uses a combination of ultrasound energy and a low thrombolytic drug dose to restore blood flow in patients with PE and other occlusions in the peripheral vasculature.

The REAL-PE study reviewed data from [Truveta](#), a data and analytics company that provides EHR data from more than 30 U.S. health systems and 100 million patients, including lab values, co-morbidities, images, demographics and clinical outcomes, as well information about the performance of specific medical devices. In the REAL-PE analysis, 2,259 patients who experienced PE and received interventional treatment with either the EKOS system or the FlowTrieve system between 2009 and 2023 were identified, and safety events associated with both devices were compared.

"In the past decade, there have been a number of advances in interventional therapies for the treatment of PE, but gaps in clinical evidence still exist when it comes to determining the optimal modality for each patient's unique needs," said Peter Monteleone, MD, FACC, FSCAI, an interventional cardiologist with Ascension and principal investigator of the REAL-PE study. "The REAL-PE study provides comprehensive data and unprecedented insight into the real-world performance of specific interventional devices, which can help physicians make more informed clinical decisions."

Major bleeding events in the REAL-PE study were based on definitions derived from established clinical criteria and guidelines, with statistically significant lower rates within seven days of the procedure consistently found in the cohort of patients treated with the EKOS system:

- A rate of 12.4% for patients treated with the EKOS system vs. 17.3% for those treated with the FlowTrieve system (p=0.0018), using the International Society on Thrombosis and Haemostasis (ISTH) definition, and; <sup>iv</sup>
- A rate of 11.8% for patients treated with the EKOS system vs. 15.4% for those treated with the FlowTrieve system (p=0.0190), using the Bleeding Academic Research Consortium type 3b (BARC3b) definition. <sup>v</sup>

Medical coding data in the analysis also demonstrated that intracerebral hemorrhage within seven days following the procedure occurred statistically significant less frequently among patients treated with the EKOS system (0.3% vs. 1.3%, p=0.005). All other studied safety events also trended in favor of the EKOS system, including in-hospital mortality (2.6% vs. 3.7%) and all-cause 30-day readmission rates (5.1% vs. 5.4%). Median lengths of hospital stay were comparable at 3.6 days for both groups.

"Electronic health record data of this scale provides in-depth information about larger, more diverse patient populations, while also accounting for multiple variables including complex medical histories or co-morbidities that often exclude patients from clinical trials," said Michael R. Jaff, D.O., chief medical officer and vice president, Clinical Affairs, Technology and Innovation, Peripheral Interventions, Boston Scientific. "While data from clinical trials and registries will always play an important role in healthcare, access to the breadth of data such as that used in the REAL-PE study has the potential to better inform and accelerate clinical decision-making and, ultimately, improve patient care."

To date, the EKOS system has been used to treat more than 100,000 patients with PE globally, and the next-generation EKOS+™ Endovascular System received U.S. Food and Drug Administration 510(k) clearance in 2022.

More information on the REAL-PE study is available [here](#) and on the EKOS system [here](#).

**About Boston Scientific**

Boston Scientific transforms lives through innovative medical solutions that improve the health of patients around the world. As a global medical technology leader for more than 40 years, we advance science for life by providing a broad range of high-performance solutions that address unmet patient needs and reduce the cost of healthcare. For more information, visit [www.bostonscientific.com](http://www.bostonscientific.com) and connect on [Twitter](#) and [Facebook](#).

### Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements within the meaning of Section 27A of the Securities Act of 1933 and Section 21E of the Securities Exchange Act of 1934. Forward-looking statements may be identified by words like "anticipate," "expect," "project," "believe," "plan," "may," "estimate," "intend" and similar words. These forward-looking statements are based on our beliefs, assumptions and estimates using information available to us at the time and are not intended to be guarantees of future events or performance. These forward-looking statements include, among other things, statements regarding our business plans and product performance and impact. If our underlying assumptions turn out to be incorrect, or if certain risks or uncertainties materialize, actual results could vary materially from the expectations and projections expressed or implied by our forward-looking statements. These factors, in some cases, have affected and in the future (together with other factors) could affect our ability to implement our business strategy and may cause actual results to differ materially from those contemplated by the statements expressed in this press release. As a result, readers are cautioned not to place undue reliance on any of our forward-looking statements.

Factors that may cause such differences include, among other things: future U.S. and global economic, political, competitive, reimbursement and regulatory conditions; new product introductions; expected procedural volumes; the closing and integration of acquisitions; demographic trends; intellectual property rights; litigation; financial market conditions; the execution and effect of our business strategy, including our cost-savings and growth initiatives; and future business decisions made by us and our competitors. New risks and uncertainties may arise from time to time and are difficult to predict. All of these factors are difficult or impossible to predict accurately and many of them are beyond our control. For a further list and description of these and other important risks and uncertainties that may affect our future operations, see Part I, Item 1A – *Risk Factors* in our most recent Annual Report on Form 10-K filed with the Securities and Exchange Commission, which we may update in Part II, Item 1A – *Risk Factors* in Quarterly Reports on Form 10-Q we have filed or will file hereafter. We disclaim any intention or obligation to publicly update or revise any forward-looking statements to reflect any change in our expectations or in events, conditions or circumstances on which those expectations may be based, or that may affect the likelihood that actual results will differ from those contained in the forward-looking statements. This cautionary statement is applicable to all forward-looking statements contained in this document.

### CONTACTS:

Blake Rouhani  
Media Relations  
+1.763.494.2268  
[Blake.Rouhani@bsci.com](mailto:Blake.Rouhani@bsci.com)

Lauren Tengler  
Investor Relations  
+1.508.683.4479  
[BSXInvestorRelations@bsci.com](mailto:BSXInvestorRelations@bsci.com)

*\*Dr. Peter Monteleone is a paid consultant of Boston Scientific. He has not been compensated in connection with this press release.*

<sup>i</sup> Centers for Disease Control and Prevention (CDC). Venous thromboembolism in adult hospitalizations -United States, 2007-2009. MMWR Morb Mortal Wkly Rep. 2012;61(22):401-404

<sup>ii</sup> Sedhom R, Megaly M, Elbadawi A, et al. Contemporary National Trends and Outcomes of Pulmonary Embolism in the United States. Am J Cardiol. 2022;176:132-138. doi:10.1016/j.amjcard.2022.03.060.

<sup>iii</sup> Giri J, Sista AK, Weinberg I, Kearon C, Kumbhani DJ, Desai ND, Piazza G, Gladwin MT, Chatterjee S, Kobayashi T, Kabrhel C, Barnes GD. Interventional Therapies for Acute Pulmonary Embolism: Current Status and Principles for the Development of Novel Evidence: A Scientific Statement From the American Heart Association. Circulation. 2019;140:e774-801.

<sup>iv</sup> Definition based on criteria from the International Society on Thrombosis and Haemostasis: Major bleeding diagnosis or a hemoglobin decrease  $\geq 2$  mg/dL and transfusion.

<sup>v</sup> Definition based on criteria from the Bleeding Academic Research Consortium: Major bleeding diagnosis or a hemoglobin decrease  $\geq 5$  mg/dL.

<https://stage.mediaroom.com/bostonscientific/2023-10-24-REAL-PE-Study-Demonstrates-Statistically-Significant-Lower-Major-Bleeding-Rates-with-the-EKOS-TM-Endovascular-System-Compared-to-Mechanical-Thrombectomy-Device-for-Treatment-of-Pulmonary-Embolism>